

# Conditional Distributions and Joint Distributions

Rob McCulloch

1. Conditional, Joint and Marginal Distributions
2. Bayes Theorem
  - 2.1. Problem: Disease Testing
  - 2.2. Problem: Classifying Defaulters
3. Several Variables
4. Independence
5. IID
  - 5.1. Problem: Making Two Parts
  - 5.2. Problem: IID Bernoulli Draws
  - 5.3. Problem: Monte Hall
  - 5.4. Problem: Democrats and Republicans, Two Draws

# 1. Conditional, Joint, and Marginal Distributions

In general we want to use probability to address problems involving more than one variable at the time.

For example we may need want to:

- ▶ describe our uncertainty about several quantities together  
(the joint distribution)
- ▶ understand how learning values about some variables affects our beliefs about others  
(the conditional distribution).

Suppose you are thinking about sales next quarter.

In order to think about sales, it may be helpful to think about sales *and* what will happen to the economy.

Let  $E$  denote the performance of the economy next quarter... for simplicity, say  $E = 1$  if the economy is expanding and  $E = 0$  if the economy is contracting (what kind of random variable is this?)

Let's assume  $p(E = 1) = 0.7$

Let  $S$  denote my sales next quarter... and let's suppose the following probability statements:

$s$	$p(S = s E = 1)$	$s$	$p(S = s E = 0)$
1	0.05	1	0.20
2	0.20	2	0.30
3	0.50	3	0.30
4	0.25	4	0.20

These are called *Conditional Distributions*

$s$	$p(S = s E = 1)$	$s$	$p(S = s E = 0)$
1	0.05	1	0.20
2	0.20	2	0.30
3	0.50	3	0.30
4	0.25	4	0.20

- ▶ In blue is the conditional distribution of  $S$  given  $E = 1$
- ▶ In red is the conditional distribution of  $S$  given  $E = 0$
- ▶ We read: *the probability of Sales of 4 ( $S = 4$ ) **given (or conditional on)** the economy is growing ( $E = 1$ ) is 0.25.*


*The way  $S$  is related to  $E$  is captured by the difference between the conditional distributions !!!*

The conditional distributions tell us about about what can happen to  $S$  for a given value of  $E$ ... but what about  $S$  and  $E$  jointly?

$$\begin{aligned} p(S = 4 \text{ and } E = 1) &= p(E = 1) \times p(S = 4|E = 1) \\ &= 0.70 \times 0.25 = 0.175 \end{aligned}$$

In English, 70% of the times the economy grows and 1/4 of those times sales equals 4... 25% of 70% is 17.5%

here is a display  
of the whole  
process:



There  
are 8 possible  
outcomes for (S,E)

We can specify the distribution of the pair of random variables  $(S, E)$  by listing all possible pairs and the corresponding probability.

$(s, e)$	$p(S = s, E = e)$
(1,1)	.035
(2,1)	.14
(3,1)	.35
(4,1)	.175
(1,0)	.06
(2,0)	.09
(3,0)	.09
(4,0)	.06

Question: What is  $Pr(S = 1)$  ?



We call the probabilities of  $E$  and  $S$  together the **joint distribution** of  $E$  and  $S$ .

In general the notation is...

- ▶  $p(Y = y, X = x)$  is the **joint probability** the random variable  $Y$  equals  $y$  **AND** the random variable  $X$  equals  $x$ .
- ▶  $p(Y = y|X = x)$  is the **conditional probability** the random variable  $Y$  takes the value  $y$  **GIVEN** that  $X$  equals  $x$ .
- ▶  $p(Y = y)$  or  $p(X = x)$  are the **marginal probabilities** of  $Y = y$  or  $X = x$

## Important relationships

Relationship between the joint and conditional...

$$\begin{aligned}Pr(Y = y, X = x) &= Pr(X = x) \times Pr(Y = y|X = x) \\ &= Pr(Y = y) \times Pr(X = x|Y = y)\end{aligned}$$

Relationship between joint and marginal...

$$\begin{aligned}Pr(X = x) &= \sum_y Pr(X = x, Y = y) \\ Pr(Y = y) &= \sum_x Pr(X = x, Y = y)\end{aligned}$$

## A Note on Notation

We have used the notations

$$P(Y = y), P(Y = y, X = x), P(Y = y | X = x)$$

You will see all kinds of similar, but not exactly the same notations for these fundamental concepts.

For example, we will sometimes use  $p(x, y)$  in place of  $P(X = x, Y = y)$  when it is clear from the context what we mean.

For example in the  $(S, E)$  example I could write  $P(S = s, E = e)$  or just  $p(s, e)$ .

## The Two-Way Table Display of the Joint Distribution

This is a nice way to display a joint distribution.

Same information as when we just listed the  $(s, e)$  pairs and their probabilities but this way we can see some things more easily.

		S				
		1	2	3	4	
E	0	.06	.09	.09	.06	.3
	1	.035	.14	.35	.175	.7
		.095	.23	.44	.235	1

For example, you can see why the marginals are called “the marginals”.

## Conditionals from Joints

We derived the joint distribution of  $(E, S)$  from the marginal for  $E$  and the conditional  $S | E$ .

You can also calculate the conditional from the joint by doing it the other way

$$Pr(Y = y, X = x) = Pr(X = x) Pr(Y = y | X = x)$$

$\Rightarrow$

$$Pr(Y = y | X = x) = \frac{Pr(Y = y, X = x)}{Pr(X = x)}$$

Example... Given  $E = 1$  what is the probability of  $S = 4$ ?

		S				
		1	2	3	4	
E	0	.06	.09	.09	.06	.3
	1	.035	.14	.35	.175	.7
		.095	.23	.44	.235	1

$$p(S = 4|E = 1) = \frac{p(S = 4, E = 1)}{p(E = 1)} = \frac{0.175}{0.7} = 0.25$$

Example... Given  $S = 4$  what is the probability of  $E = 1$ ?

		S				
		1	2	3	4	
E	0	.06	.09	.09	.06	.3
	1	.035	.14	.35	.175	.7
		<hr/>				
		.095	.23	.44	.235	1

$$p(E = 1|S = 4) = \frac{p(S = 4, E = 1)}{p(S = 4)} = \frac{0.175}{0.235} = 0.745$$

## 2. Bayes Theorem

So, in general, you can compute the joint from marginals and conditionals and the other way around.

How you think about stuff depends on what's easiest or what you know, or what you care about.

Suppose you toss two coins:  $X$  is the first,  $Y$  is the second.

In each case 1 means a head and 0 a tail.

What is  $P(X = 1, Y = 1) = P(\text{two heads})$  ?

(1) Directly figure out the joint distribution.

There are 4 possible outcomes for the two coins and each is equally likely so it is  $1/4$ .

(2) Figure out some marginals and conditionals.

$$P(X = 1, Y = 1) = P(X = 1) * P(Y = 1 | X = 1) = (1/2) * (1/2) = 1/4.$$



*Bayes Theorem* refers to the situation where we build a model for  $(Y, X)$  by thinking about

$$Pr(Y = y | X = x), Pr(X = x).$$

and then, having observed  $Y = y$  compute

$$Pr(X = x | Y = y)$$


## Example: Disease Testing

Disease testing example...

Let  $D = 1$  indicate you have a disease.

Let  $T = 1$  indicate that you test positive for it

We know the marginal of  $D$  and the conditional of  $T$  given  $D$ .



If you take the test and the result is positive, you are really interested in the question: **Given that you tested positive, what is the chance you have the disease?**

		D	
		0	1
T	0	.9702	.001
	1	.0098	.019

$$p(D = 1 | T = 1) = \frac{0.019}{(0.019 + 0.0098)} = 0.66$$

The computation of  $p(x|y)$  from  $p(x)$  and  $p(y|x)$  is called Bayes theorem...

$$p(x|y) = \frac{p(y, x)}{p(y)} = \frac{p(y, x)}{\sum_x p(y, x)} = \frac{p(x)p(y|x)}{\sum_x p(x)p(y|x)}$$

In the disease testing example:

$$p(D = 1 | T = 1) = \frac{p(T=1|D=1)p(D=1)}{p(T=1|D=1)p(D=1) + p(T=1|D=0)p(D=0)}$$

$$p(D = 1 | T = 1) = \frac{0.019}{(0.019+0.0098)} = 0.66$$

## 2.1. Problem: Disease Testing

Suppose a person is randomly drawn from a large population and then tested for a disease.

Let  $D = 1$  if the person has the disease and 0 otherwise.

Let  $T = 1$  if the person tests positive and 0 otherwise.

Suppose

$$P(D = 0) = .99.$$

$$P(T = 1 \mid D = 0) = .01.$$

$$P(T = 1 \mid D = 1) = .97.$$

(a)

Draw the diagram depicting the marginal of  $D$  and the conditional of  $T \mid D$ .

(you know, the one that branches as you go left to right).

(b)

Give the joint distribution of  $D$  and  $T$  in the two way table format.

(c)

What is  $P(D = 1 \mid T = 1)$ ?

## Solution

(b)

	T0	T1
D0	0.9801	0.0099
D1	0.0003	0.0097

```
> .99*.99
```

```
[1] 0.9801
```

```
> .99*.01
```

```
[1] 0.0099
```

```
> .01*.03
```

```
[1] 3e-04
```

```
> .01*.97
```

```
[1] 0.0097
```

(c)

$$P(D = 1 | T = 1) = P(D = 1, T = 1) / P(T = 1) = .0097 / (.0097 + .0099) = 0.494898.$$

## 2.2. Problem: Classifying Defaulters

Let's suppose a Credit Card company defines 4 "risk classes" based on the outstanding balance on the card.

risk class 1: balance in  $(0,500]$

risk class 2: balance in  $(500,1000]$

risk class 3: balance in  $(1000,1500]$

risk class 4: balance in  $(1500, \infty)$  (that is, anything  $> 1500$ )

Let  $C$  denote the random variable denoting the risk class of an account and  $D$  denote the random Variable which is 1 if there is a default and 0 if not. The research department has come up with the conditional distribution of  $C$  for  $D=0$  or 1.

	$p(c D=0)$	$p(c D=1)$
1	0.27	0.00
2	0.39	0.02
3	0.27	0.21
4	0.07	0.77

(a)

Suppose  $P(D = 1) = .03$ .

Compute  $P(D = 1|C = c)$  for  $c = 1, 2, 3, 4$ .

(b)

Suppose  $P(D = 1) = .5$ .

Compute  $P(D = 1|C = c)$  for  $c = 1, 2, 3, 4$ .



## Solution

(a)

If  $P(D = 1) = .03$  then the joint distribution of  $(C, D)$  is given by the table:

	d=0	d=1
c=1	0.2619	0.0000
c=2	0.3783	0.0006
c=3	0.2619	0.0063
c=4	0.0679	0.0231

For example

$$p(C = 4, D = 0) = p(D = 0)p(D = 4|D = 0) = .97*.07 = 0.0679.$$

To get the conditional probs  $p(D = 1|c)$  divide the second column by the sum of the two columns:

	c=1	c=2	c=3	c=4
	0.000000000	0.001583531	0.023489933	0.253846154

For example,

$$p(D = 1|C = 2) = .0006/ (.0006 + .3783) = 0.001583531.$$

(b)

If  $P(D = 1) = .5$  then the joint distribution of  $(C, D)$  is given by the table:

	d=0	d=1
c=1	0.135	0.000
c=2	0.195	0.010
c=3	0.135	0.105
c=4	0.035	0.385

For example  $.01 = .5 * .02$

$$P(C = 2, D = 1) = P(D = 1) * P(C = 2|D = 1).$$

And the conditional  $P(D = 1|c)$  is:

	c=1	c=2	c=3	c=4
	0.0000000	0.04878049	0.43750000	0.91666667

For example,  $.01 / (.01 + .195) = 0.04878049$ .

The marginals  $P(D = 1)$  is called the *prior probability* and the conditional  $P(D = 1|c)$  is called the *posterior probability*.

Clearly, the prior makes a difference!!!

### 3. Several Variables

Of course, we often want to think about more than two variables at a time.

We can extend the ideas we used with two variables to many variables.

Suppose we have the three random variables,

$$(Y_1, Y_2, Y_3)$$

Then,

$$Pr(Y_1 = y_1, Y_2 = y_2, Y_3 = y_3) =$$

$$Pr(Y_1 = y_1) Pr(Y_2 = y_2 \mid Y_1 = y_1) Pr(Y_3 = y_3 \mid Y_1 = y_1, Y_2 = y_2)$$

Or, using more succinct notation:

$$p(y_1, y_2, y_3) = p(y_1) p(y_2 | y_1) p(y_3 | y_1, y_2)$$

*You can keep going for any number of variables !!*

## Example

Suppose we have 10 voters.

4 are republican and 6 are democratic.

We randomly choose 3.

Let  $Y_i$  be 1 if the  $i^{\text{th}}$  voter is a democrat and 0 otherwise,  
 $i = 1, 2, 3$ .

What is

$$\Pr(Y_1 = 1, Y_2 = 1, Y_3 = 1)$$

What is the probability of getting three democrats in a row ??

$$Pr(Y_1 = 1, Y_2 = 1, Y_3 = 1) =$$

$$Pr(Y_1 = 1) p(Y_2 = 1 | Y_1 = 1) Pr(Y_3 = 1 | Y_1 = 1, Y_2 = 1)$$

$$= (6/10)(5/9)(4/8)$$

$$= (1/6) = .167.$$

When we randomly pick a person from a population of people, and then randomly pick a second from the ones left, and so on, we call it *sampling without replacement*.

If we put the person back each time and randomly choose from the whole group each time, then we call it *sampling with replacement*.

Suppose we sample from our 10 voters with replacement.

Now what is

$$\Pr(Y_1 = 1, Y_2 = 1, Y_3 = 1)$$



$$Pr(Y_1 = 1, Y_2 = 1, Y_3 = 1) =$$

$$Pr(Y_1 = 1) p(Y_2 = 1 | Y_1 = 1) Pr(Y_3 = 1 | Y_1 = 1, Y_2 = 1)$$

$$= (6/10)(6/10)(6/10)$$

$$= .6^3 = .216$$

Notice that when we sample with replacement

$$p(Y_2 = 1 \mid Y_1 = y_1) \text{ and } Pr(Y_3 = 1 \mid Y_1 = y_1, Y_2 = y_2)$$

do not depend on  $y_1$  and  $y_2$ .

What happens for  $Y_1$  does not affect what we think will happen for  $Y_2$  and what happens for  $Y_1$  and  $Y_2$  does not affect what will happen for  $Y_3$ .

In this case we say the random variables are *independent*.

## 4. Independence

Given a bunch of random variables, we say they are independent of each other if the conditional distribution of any one of them does not depend on anything you might observe for any of the others.

### Example

Suppose I am about to toss 100 coins.

Let  $Y_i$  be 1 if the  $i^{\text{th}}$  coin is a head and 0 otherwise.

What is  $Pr(Y_3 = 1)$ ?

What is  $Pr(Y_3 = 1 \mid Y_1 = 1, Y_2 = 0)$  ?

What is  $Pr(Y_3 = 1 \mid Y_1 = 0, Y_2 = 1)$  ?

What is  $Pr(Y_3 = 1 \mid Y_1 = 0, Y_2 = 0)$  ?

What is  $Pr(Y_3 = 1 \mid Y_1 = 1, Y_2 = 1)$  ?

What is

$Pr(Y_{100} = 1 \mid Y_1 = 1, Y_2 = 1, \dots, Y_{99} = 1)$  first 99 are heads?

What is

$Pr(Y_1 = 1, Y_2 = 1, Y_3 = 1, \dots, Y_{100} = 1)$  100 heads in a row!?!?

## Independence, Conditional Equals Marginal

If random variables are independent then the conditional is the marginal.

For two random variables  $X$  and  $Y$  if  $X$  and  $Y$  are independent then,

$$Pr(Y = y | X = x)$$

does not depend on  $x$ .

We also have:

$$Pr(Y = y | X = x) = Pr(Y = y)$$

*What you believe about  $Y$  knowing  $X = x$ , is the same as what you believe if you know nothing about  $X$ .*

## Example

$X$  is 1 if first coin is head.

$Y$  is 1 if second coin is head.

What is

$$P(Y = 1 \mid X = 1)?$$

What is

$$P(Y = 1 \mid X = 0)?$$

What is

$$P(Y = 1)?$$

## Independence, Joints, and Marginals

If  $X$  and  $Y$  are independent then the joint is the product of the marginals:

$$\begin{aligned} p(x, y) &= p(x) p(y | x) \\ &= p(x) p(y) \end{aligned}$$

This also works “the other way”, that is, if the joint is the product of the marginals then they are independent.

## Example

You are about to manufacture two parts.

$X = 1$  if part one fails, 0 else.

$Y = 1$  if part two fails, 0 else.

The table below gives the joint distribution of  $X$  and  $Y$ .

		X	
		0	1
Y	0	.72	.08
	1	.18	.02



		X	
		0	1
Y	0	.72	.08
	1	.18	.02

$$Pr(Y = 1 | X = 0) = .18 / .9 = .2$$

$$Pr(Y = 1 | X = 1) = .02 / .1 = .2$$

$$Pr(Y = 1) = .18 + .02 = .2$$

$$Pr(Y = 1, X = 1) = .02$$

$$Pr(Y = 1)Pr(X = 1) = .2 * .1 = .02.$$

X and Y are independent.

For random variables  $Y_i$ , if they are independent we have

$$\begin{aligned} p(y_1, y_2, \dots, y_n) &= \\ p(y_1) p(y_2 | y_1) p(y_3 | y_1, y_2) \dots p(y_n | y_1, y_2, \dots, y_{n-1}) \\ &= p(y_1) p(y_2) p(y_3) \dots p(y_n) \end{aligned}$$

### Example

If  $Y_i$  is 1 if the  $i^{\text{th}}$  coin is a head, 0 else,  $i = 1, 2, \dots, 10$ , what is

$$p(1, 1, \dots, 1)$$

10 heads in a row?

$$\begin{aligned} p(1, 1, \dots, 1) &= \\ &= p(1) p(1) p(1) \dots p(1) \\ &= .5^{10} = 0.0009765625. \end{aligned}$$

## 5. IID

Suppose we are about to toss 100 coins.

Let  $Y_i$  be 1 if heads, 0 else,

We usually think the  $Y$ 's are independent.

In addition, we usually think they are *identically distributed*, that is, each one has *the same marginal distribution*.

What is  $Pr(Y_{20} = 1)$ ?

What is  $Pr(Y_{98} = 1)$ ?

When random variables are independent and identically distributed we say they are **IID**.

**I**: independent

**ID**: identically distributed.

In our coins example, each  $Y_i \sim \text{Bernoulli}(.5)$ .

We can succinctly describe coin tossing by

$$Y_i \sim \text{Bernoulli}(.5), \text{ IID}$$

## Example

Suppose we have 10 voters. 4 are republican and 6 are democratic.

We randomly choose 3, sampling *with replacement*.

Let  $Y_i$  be 1 if the  $i^{\text{th}}$  voter is a democrat and 0 otherwise,  $i = 1, 2, 3$ .

How can we describe the joint distribution of  $(Y_1, Y_2, Y_3)$ , are they IID?

## Example

Suppose we have 10 voters. 4 are republican and 6 are democratic.

We randomly choose 3, sampling *without replacement*.

Let  $Y_i$  be 1 if the  $i$  th voter is a democrat and 0 otherwise,  
 $i = 1, 2, 3$ .

How can we describe the joint distribution of  $(Y_1, Y_2, Y_3)$ , are they IID?

## Example

Suppose we have 1,000,000 voters. 400,000 are republican and 600,000 are democratic.

We randomly choose 3, sampling *without replacement*.

Let  $Y_i$  be 1 if the  $i^{th}$  voter is a democrat and 0 otherwise,  $i = 1, 2, 3$ .

How can we describe the joint distribution of  $(Y_1, Y_2, Y_3)$ , are they IID?



## Example

Suppose I am about to toss a die 100 times.

Let  $D_i$  be the outcome for the  $i^{\text{th}}$  toss  
(a number in  $\{1, 2, 3, 4, 5, 6\}$ ).

Are the  $D_i$  IID?

## Example

Suppose an experienced NBA player is about to take repeated free-throws.

Let  $Y_i$  be 1 if he makes the  $i^{\text{th}}$  attempt and 0 otherwise.

Are these  $Y_i$  iid Bernoulli?

This is known as the “hot hand” question.

Most people who play sports believe that they can get “hot” so that if they made the last few, they are more likely to make the next one.

However, if you look at the data, it looks IID Bernoulli!!

How do you look at the data to see if it looks IID Bernoulli. That is covered in the notes “Modeling with IID Bernoulli Draws”.

## Example

Suppose the first penalty in an NHL game is on team A.

For subsequent penalties  $P_i = 1$  if the penalty is on a different team than the previous one and 0 otherwise.

Are the  $P$ 's independent?

Are they IID?

These are not IID.

If the last two (or three!) penalties were on the same team, it becomes quite a bit more likely that the next penalty will be on the other team.

See

Reversal of fortune: a statistical analysis of penalty calls in the national hockey league", (2014), Journal of Quantitative Analysis in Sports 10 (2), 207-224 (Jason Abrevaya and Robert McCulloch)

## Example

Suppose you are monitoring a stock and for every 10 minute interval, you record whether the price went up or down.

Let  $U_i$  be 1 if it goes up in the  $i^{\text{th}}$  interval, 0 otherwise.

Are the  $U_i$  IID?


Of course, this is a much studied question.

We leave this to your finance courses but just note that it is very interesting how little dependence there is!!!!


## Example: Modeling Defects:

You are in charge of a manufacturing process.

For the last 300 parts made, we have 1 if the part is defective and 0 otherwise.



$54/300 = .18$  of  
the parts are de-  
fective.






We are about to make the next part.

Let  $Y$  be 1 if it turns out to be defective and 0 if it is good.


What is the distribution of  $Y$ ?

Here are 300 coin tosses, 1 if heads, 0 if tails.




$142/300 = .47$   
of the coins are  
heads.

If  $Y = 1$  means  
the next one  
turns out to be  
a head, what is  
the distribution  
of  $Y$ ?





Here are 300 observations on another 0/1 process.  
The second plot just shows the last 100.



$60/300 = .20$  of the observations are 1.

If  $Y$  denotes the next one, what is the distribution of  $Y$ ?





The defects look like the kind of thing you would get as IID draws from a Bernoulli, with  $p = .18$ .

We model:

$$Y_i \sim \text{Bernoulli}(.18).$$

So, a plausible choice for distribution of the next one is just  $Y \sim \text{Bernoulli}(.18)$ .

What is the probability the next part made is defective.?

What is the probability the next 10 parts are good?

## Note:

For  $Y_i \sim \text{Bernoulli}(p)$ , the usual estimator of  $p$  is the observed fraction of 1's.

If we call this estimator  $\hat{p}$ , then,

- ▶ we have the *standard error*:  $se(\hat{p}) = \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$ .
- ▶ and the associated 95% *confidence interval* for  $p$ :  $\hat{p} \pm 2se(\hat{p})$ .

## Defect Example:

$$\hat{p} = .18, se(\hat{p}) = \text{sqrt}(.18*(1-.18)/300) = .022$$

Approximate 95% confidence interval:  $.18 \pm .044 = (.14, .22)$ .

Is it big?

## Note:

For the the “sampling from a finite population problem” and the defects we used the IID Bernoulli model.

*But,*

*the way we justify the use of the model  
is completely different !!!!!*

## Note:

For the deterministic 0/1 process we can summarize the numbers by saying we observed 20% ones. But, is that an estimate of anything?

## 5.1. Problem: Making Two Parts

You are about to make two parts.

Let  $X = 1$  if the first part is defective and 0 otherwise.

Let  $Y = 1$  if the second part is defective and 0 otherwise.

The table below gives the joint probability distribution for the two random variables  $X$  and  $Y$ .

	X0	X1
Y0	0.81	0.09
Y1	0.09	0.01

- (a) What is the marginal distribution of  $X$ ?
- (b) What is the marginal distribution of  $Y$ ?
- (c) What is the conditional distribution of  $Y \mid X = 1$ ?
- (d) What is the conditional distribution of  $Y \mid X = 0$ ?
- (e) Are  $X$  and  $Y$  independent?
- (f) Is  $p(x, y) = p(x) p(y)$  for  $x = 0, 1, y = 0, 1$ .
- (g) Are  $X$  and  $Y$  IID?
- (h) What are the mean and variance of  $X$ ?



## Solution

(a)  $P(X = 0) = .9, P(X = 1) = .1.$

Or,  $X \sim \text{Bernoulli}(.1).$

Or,

x	p(x)
0	.9
1	.1

(b) Same as  $X.$

(c) Same as marginal for  $Y.$

(d) Same as marginal for  $Y.$

(e) They are independent because all conditionals are the same as the marginal.

(f) Yes, because they are independent.

(g) Yes. They are both  $\text{Bernoulli}(.1).$

(h)  $E(X) = .1, \text{Var}(X) = .1 * .9 = .09.$

## 5.2. Problem: IID Bernoulli Draws

Computers can flip coins!!

A very important part of modern computation is simulation.

We can ask the computer of IID draws from a distribution.

Let's ask the computer for 100 IID draws,

$$X_i \sim \text{Bernoulli}(.5), \quad i = 1, 2, \dots, 100.$$

In excel:

To get iid Bernoulli draws:

/data/Data Analysis/Random Number Generation:

Number of variables: 1

Number of random Numbers: 100

Distribution: Bernoulli

p-value: .5

Random Seed: 99

output range: a2:a101

Then put the name x in a1.

To get a time series plot of the draws:

click on a cell containing one of the draws.

/insert/line

play around with different choices of line type.

## In R:


```
##to get iid Bernoulli draws
set.seed(99)
x = rbinom(100,1,.5) #100 iid draws ~ Bernoulli(.5)

##to get a time series plot of x:
plot(x,type="b") #type="b" gives lines and points
```

- (a) Do the time series plot of your IID draws.  
(time series plot is just  $y_1, y_2 \dots$  v.s.  $1, 2, \dots$ )
- (b) What is the average value of the draws?
- (c) Do 5 draws twice, each time compute the average.
- (d) Do 100 draws twice, each time compute the average.
- (e) How are (c) and (d) different?
- (f) Get 100 iid Bernoulli(.1) draws. What is the average of the draws?

## Solution

(a) Here is what I got.



Of course, you will get something different.

(b) I got .46. (c) I got 0 and .2.

(d) I got .5 and .53.

(e) With 100 draws, both averages are pretty close to  $p=.5$ , not so with 5!

(f) I got .08, pretty close to  $p = .1$ .

## 5.3. Problem: Monte Hall

This problem is named after the host of the long running TV show *Let's make a deal*. There has been a vigorous debate about what the correct answer is!!

A contestant must choose one of three closed doors. There is a prize (say a car) behind one of the three doors. Behind the other two doors, there is something worthless (traditionally a goat). After the contestant chooses one of the three doors, Monty opens one of the other two, revealing a goat (never the car!!). There are now two closed doors. The contestant is asked whether he would like to switch from the door he initially chose, to the other closed door.

The contestant will get whatever is behind the door he has finally chosen. Should he switch?

# Solution

Assume (I claim without loss of generality) that you initially select door 1.

The game is about to be played.


Up to the point where you must decide whether or not to switch, there are two things about which we are uncertain:

C: the door the car is behind, C is 1, 2, or 3

M: which door Monte will open, M is 2 or 3 (given you have selected door 1).

We need the joint distribution of (C,M).

We will first write it in terms of the marginal for C and the conditional for M given C since it is the most obvious in this form.





## 5.4. Problem: Democrats and Republicans, Two Draws

Suppose we have 5 Democrats and 5 Republicans.

We are about to sample two voters.

Let  $Y_1$  be 1 if the first voter is a dem and 0 else.

Let  $Y_2$  be 1 if the second voter is a dem and 0 else.

(a)

Suppose we sample without replacement.

Give the 2x2 table representing the joint distribution of  $Y_1$  and  $Y_2$ .

(b)

In (a), are the two  $Y$ 's independent?

Are they identically distributed?

Are they IID?

(c)

Now suppose we sample the two voters with replacement.  
Give the 2x2 table representing the joint distribution of  $Y_1$  and  $Y_2$ .

(d)

In (c), are the two  $Y$ 's independent?

Are they identically distributed?

Are they IID?

(e)

Suppose we had 5 million of each instead of just 5.

How would your answers to (a)-(d) change?

## Solution

(a)

	Y1=0	Y1=1
Y2=0	0.222	0.278
Y2=1	0.278	0.222

note:  $(5/10)*(4/9) = .222$  (after rounding).

$(5/10)*(5/9) = .278$  (after rounding).

(b)

They are not independent.

They are identically distributed.

They are not IID.

(c)

	Y1=0	Y1=1
Y2=0	0.25	0.25
Y2=1	0.25	0.25

(d)

Yes,  $Y_1, Y_2$  are iid Bernoulli(.5).

(e)

Now our joint in (c) is basically the same as in (a) so in both cases the  $Y$ 's are iid Bernoulli(.5).