

Modeling with IID Normal Draws

Homework Problems

Rob McCulloch

1. The Histogram and IID Draws
2. The Normal Distribution and Data
3. Standardization
 - 3.1. Problem: Australian Return Data
 - 3.2. Problem: Unusual Belgian Return?
 - 3.3. Problem: The Random Walk

3.1. Problem: Australian Return Data

(a)

Do the time series plot and histogram of the “australia” returns from the countries data (conret.csv). These are returns on a portfolio of Australian equities (stocks).

Does it look reasonable to model the returns as iid draws from a normal distribution?

Note: the “time series plot” is just i versus x_i for values x_i .

The is also called the sequence plot. We have already used this plot a lot!!

In Excel: Select the column of Australian returns and then go to /Insert and then play around to find the plot type you like the best.

If you just do scatterplot, it seems to look ok.

In R:

```
cdat = read.csv("http://www.rob-mcculloch.org/data/conret.csv")
plot(cdat$australia,type="b",xlab="year number",
      ylab="Australia Returns",col="blue")
```

(b)

Let μ be the sample average of the australian returns and σ be the sample standard deviation.

If we go ahead and model our returns as iid normal and use these values for μ and σ , what is the probability of a negative return?

(c)

What is the “z” value corresponding to $r = 0$

($z = (r - \mu)/\sigma$)?

Check that $P(Z < z)$ gives you the same number you got in part (b), where $Z \sim N(0, 1)$.

(d)

z each australian return.

Do the histogram of the z values.

Does it kind of look like the standard normal pdf?!!

In Excel you could copy the formula

`= (B2 - AVERAGE(B2:B108)) / STDEV.S(B2:B108)`

in cells 2:108 in column Y (or any empty column).

Remember to copy a formula:

- ▶ click in the top cell (e.g. Y2))
- ▶ shift/click in the bottom cell (e.g. Y108))
- ▶ type in the formula
- ▶ enter Cntl/return.

In R:

```
za = (cdat$australia - mean(cdat$australia)) / sd(cdat$australia)
```

3.2. Problem: Unusual Belgian Return?

Do the time series plot of the “belgium” returns form conret.csv.

Notice that the first return seems like an “outlier”.

Let μ be the sample average of the returns, excluding the first one.
Let σ be the sample standard deviation of the returns, excluding the first one.

What is the z value for the first return?

Is it more unusual than Gretzky?

3.3. Problem: The Random Walk Model

Get the data Price-level.csv from the webpage.

There is just one column called Price.

The data are the monthly price of an asset for 500 months (about 41 years).

(a)

Do a time series plot of the prices.

Could they be modeled as iid ?

(b)

Compute the price differences.

If the prices are denoted by P_t , then the differences are

$$D_t = P_t - P_{t-1}, t = 2, 3, \dots, n.$$

For example, the first three prices are 0.000000, -1.748190, and 4.707081

so the first two differences are

$$-1.748190 - 0.000000 = -1.74819 \text{ and}$$

$$4.707081 - (-1.748190) = 6.455271$$

```
> Price[1:4]
[1] 0.000000 -1.748190 4.707081 8.905969
> -1.748190-0.000000
[1] -1.74819
> 4.707081 - (-1.748190)
[1] 6.455271
```

Note that in R we could get the differences by

```
> ii = 2:n
> D = Price[ii]-Price[ii-1]
> print(D[1:3])
[1] -1.748190 6.455271 4.198888
```


Do a time-series plot and a histogram of the differences.

Do the differences look iid normal?

(c)

Give a distribution describing what the next price difference will be.

That is, our last observed price is $P_{500} = 609.298161$.

So the next difference is $P_{501} - 609.298161$ which is a number we are unsure about because we don't know the next price P_{501} .

(d)

Give a distribution describing what the next price (P_{501}) will be.

Our model is:

$$D_t = P_t - P_{t-1} \sim N(\mu, \sigma^2).$$

This is the very famous *random walk model* which we can express as

$$P_t = P_{t-1} + D_t$$

The next price, is the current price plus a “random” increment. In this problem we modeled the increment as iid normal.